

**Label****REVOLYTE**

Complementary Feed  
Vitamin, Mineral and Amino Acid Solution

**COMPOSITION**

1 mL solution contains:

Dietary additives			
3a820	(Thiamine hydrochloride)	Vitamin B <sub>1</sub>	0.10 mg
3a	(Riboflavin sodium phosphate)	Vitamin B <sub>2</sub>	0.04 mg
3a831	(Pyridoxine hydrochloride)	Vitamin B <sub>6</sub>	0.10 mg
3a	(Cyanocobalamin)	Vitamin B <sub>12</sub>	0.05 µg
3a315	(Niacinamide)	Vitamin B <sub>3</sub>	1.5 mg
3a842	(D-panthenol)	Vitamin B <sub>5</sub>	0.05 mg
3c321	(L-Lysine monohydrochloride)	Lysine	0.03 mg
3c301	DL-methionine		0.01 mg
2b17018	L-phenylalanine		0.03 mg
3c410	L-threonine		0.02 mg
3c440	L-tryptophan		0.01 mg
3c370	L-valine		0.05 mg
3c361	L-arginine		0.025mg
2b17033	L-cysteine		0.01 mg
2b17008	L-histidine		0.01 mg
2b17010	DL-isoleucine		0.01 mg
2b17012	L-leucine		0.04 mg
2b621	Monosodium glutamate		0.04 mg
2b	Phenol		0.1 mg
11.1.6	(Calcium chloride)	Calcium	0.23 mg
11.2.2	(Magnesium sulphate heptahydrate)	Magnesium	0.29 mg
11.5.1	(Potassium chloride)	Potassium	0.20 mg
13.2.2	Dextrose		45.46 mg
Technological additives- preservatives (1a) 1.0 mg, Carrier- water			

**ANALYTICAL CONSTITUENTS**

Moisture content 98.7%, Sugar 4.5%, Calcium 0.008%, Magnesium 0.006%, Potassium 0.036%, Sodium 0.0007%, Crude fat 0.0%, Crude protein 0.02%, Crude fiber 0.2%, Crude ash 0.75 %, Phosphorus 0.0%, Lysine 0.002%, Methionine 0.001%

**DIRECTIONS FOR USE**

Recommended to be used as a supportive and preventative supplement in cases of dehydration, electrolyte imbalance and hypoproteinemia which can occur in post-operations, blood loss, diarrhea, shock and exhaustion. Can also be used during period of convalescence.

Produced as sterile (E.Ph.) of a high quality.

<b>Animal Species</b>	<b>Recommended Usage</b>
Cattle, camels, horses, swine	500 – 1000 mL (1-2 pieces of product)
Calves, sheep, piglets	100 – 200 mL
Cats, dogs	50 – 100 mL

The product may be given directly to individual animals and to groups of animals via the feed or drinking water to supplement the daily ration to suit the required needs. Advised dosages are for 1-2 L of water or 1-2 kg of feed. Recommended usage for poultry is 500 mL product to 500 L of drinking water. It is advised to give to chicks before transportation for prevention of losses.

## **GENERAL WARNINGS**

Consult your veterinarian before use.

If the additive is administered via water for drinking protein excess should be avoided

**Net volume:** 500 mL and 1000 mL

**Shelf life:** 3 years from date of production.

## **SPECIAL PRECAUTIONS FOR STORAGE AND TRANSPORTATION**

Protect from direct sunlight. Do not freeze. Store at room temperature 15-25 °C and in a dry place. Keep out of reach of children.

## **SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIALS DERIVED FROM THE USAGE**

Do not empty into drains or with the household waste. The current legal regulations should be observed. This will help protect the environment.

## **MANUFACTURER**

FERAMED BV

Barneveld- The Netherlands

**αNL 1014**

**Batch No:**

**Man. Date:**

**Expiry Date:**